

AD-A270 936



DOCUMENTATION PAGE

Form Approved
OMB No 0704 0188

Estimate is based on average 1-hour per response, including the time for reviewing instructions, searching existing data sources, gathering and reviewing the collected information, sending comments regarding this burden estimate or any other aspect of this burden estimate, including this burden estimate, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

2. REPORT DATE
June 8, 19933. REPORT TYPE AND DATES COVERED
Final 1 May 88-30 Apr 93

4. TITLE AND SUBTITLE

THE ROLE OF AXON-SCHWANN CELL INTERACTIONS IN NERVOUS
SYSTEM IONIC HOMEOSTASIS

6. AUTHOR(S)

Edward M. Lieberman

DTIC
ELECTE

OCT 20 1993

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

East Carolina University
School of Medicine
Greenville, NC 27858-4354

5. FUNDING NUMBERS

DAALO3-88-K-0102

(2)

8. PERFORMING ORGANIZATION
REPORT NUMBER

9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)

U.S. Army Research Office
P. O. Box 12211
Research Triangle Park, NC 27709-221110. SPONSORING/MONITORING
AGENCY REPORT NUMBER

ARO 25752.10-LS

11. SUPPLEMENTARY NOTES

The view, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.

12a. DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

The long term goal of the research program of this laboratory is to reach an understanding of the physiological interactions between neurons and their associated glia that relate to their ability to cooperatively regulate the ionic and neurohumoral milieu of the perineural and/or periaxonal space. To this end we have been investigating the mechanisms by which axons and neurons generate the chemical signals during their excitation that signals the glia to activate their metabolic and uptake processes that inactivate neurotransmitters preventing their accumulation to toxic concentrations and the transport of potassium and of sodium to maintain appropriate amounts of each ion in the perineural space thereby preserving excitability properties of the neural elements of the nervous system. For these investigations we have used the intact nerve fiber - Schwann cell preparations of crayfish and squid to study nerve-Schwann cell interactions in intact systems and the mammalian Schwann cell culture to determine the generality of our findings to mammalian systems.

14. SUBJECT TERMS

Schwann cell, glia, nerve, potassium homeostasis, glutamate,
acetylcholine, nerve-glia interaction

15. NUMBER OF PAGES

16. PRICE CODE

17. SECURITY CLASSIFICATION
OF REPORT
UNCLASSIFIED18. SECURITY CLASSIFICATION
OF THIS PAGE
UNCLASSIFIED19. SECURITY CLASSIFICATION
OF ABSTRACT
UNCLASSIFIED20. LIMITATION OF ABSTRACT
UL

**Best
Available
Copy**

Statement of Problem:

In recent years it has been recognized that glia of the nervous system participate in the acute management of the ionic and neurohumoral milieu of the neural microenvironment. It is the glia that modulate changes in the ionic environment of nerve fibers and synaptic regions of the nervous system may influence excitability properties of nerve cells, release of transmitter, propagation velocity and safety factors for excitation and branch point action potential conduction. The questions addressed by this program of research are (1) what are the specific roles of the glia in neural microenvironment homeostasis (2) what mechanisms of communication exist between neurons and glia and (3) what changes in physiological properties of glia are induced by neural activity.

Our findings since the beginning of this project can be summarized as follows:

1. Glutamate and potassium released by the excited nerve fiber acts as the signal to the surrounding glia.
2. The release of glutamate and potassium appear to be coupled so that the quantity of glutamate released is in direct proportion to the potassium released thus providing an appropriate signal to the glial cell for the rapid clearance of the excess potassium that tends to appear in the perineural space during excitation.
3. The Schwann cell (glial cell) has, on its membrane, a specific non-MNDA (quisqualate/kainate) type glutamate which activates 2 separate processes.
 - a. opening of a sodium ion channel that tends to depolarize the membrane potential of the Schwann cell. The function of this particular channel is unknown but we have speculated that the sodium that enters the glial cell under these circumstances insures that the sodium/potassium transporter is primed for potassium uptake processes by the sodium pump.
 - b. activation of an increase in intracellular ionic calcium via an inositol phosphate second messenger that is required for the release of acetylcholine from the Schwann cell. ACh acts on an Schwann cell autoreceptor, coupled to a c-AMP generating mechanism that is ultimately responsible for a chloride-dependent Schwann cell hyperpolarization.
4. The Schwann cell hyperpolarization, because it is due to a decrease in chloride permeability, decreases the efflux of potassium from the Schwann cell. This allows an increase in nerve potassium efflux to an amount equivalent to the decrease in glial potassium efflux without increasing perineural potassium concentration. When excessive activity tends to cause accumulation of potassium both sodium/potassium/2chloride cotransport activity and sodium/potassium antiport transport are also activated in the glial cell to insure that the perineural microenvironment concentration of potassium remains consistent with preservation of neural excitability.
5. Enzymes of the glutamate-glutamine neurotransmitter cycle have been

93-24830



589

93 10 18 194

shown to exist in the intact nerve -glial cell preparation used in these studies as it has been identified in mammalian nervous systems. That is, glutamine synthetase, the enzyme required for conversion of glutamate to glutamine is almost exclusively compartmentalized to the glial cell while high concentrations of glutaminase are found in both glia and nerve. These findings are consistent with the concept that the glia have a primary role in removing neuroactive substances from the neural microenvironment before they accumulate to toxic levels. The presence of glutaminase in both cells is consistent with the metabolic importance of glutamate in general cell metabolism of all cells.

6. In keeping with the strong physiological coupling between nerve cells and their associated glia and the large transport activity of the glia we have also determined that approximately 90% of the metabolic activity of intact nerve tissue can be accounted for by glial cells and that the level of activity at any time is dependent on the metabolic state of the nerve cells.
7. In cultured mammalian Schwann cells we have determined that the normal resting membrane potential of these cells is low (about -45 mV) and similar to the resting potential of intact glia of the invertebrate nervous systems we have been using for *in vivo* studies.
8. In mammalian Schwann cells there is a strong coupling between the activity of the sodium/potassium/2 chloride cotransporter and the sodium pump. Increasing c-AMP of the Schwann cell activates both transporter systems. A long term requirement for high sodium pump activity results in an increase in its activity through the process of translocation of sequestered intracellular pump molecules to the plasma membrane.
9. Most recently we have been examining the manner in which mammalian glia deal with proton transport and have found them to be active pH buffers for extracellular space in addition to the requirement to maintain intracellular pH for intracellular enzymatic activity.

Publication directly attributed to support under this grant project

1. Hargittai, P.T., D.D. Ginty and E.M. Lieberman. A pyrene fluorescence technique and microchamber for measurement of oxygen consumption of single isolated axons. *Analytical Biochemistry* **163**: 418-426, 1987.
2. Abbott, N.J., E.M. Lieberman, Y. Pichon and Y. Larmet. Periaxonal K^+ regulation in the small squid *Alloteuthis*: studies on isolated and in situ axons. *Biophysical J.* **53**:275-279 (1988)
3. Brunder, D.G. and E.M. Lieberman. Studies of axon-glia interactions and periaxonal K^+ homeostasis. I. The influence of K^+ , Cl^- and cholinergic agents on the membrane potential of the adaxonal glia of the crayfish medial giant axons. *Neuroscience* **53**(3) 951-959 (1988)
4. S. Hassan and Lieberman, E.M. Studies on axon-glia interactions and periaxonal K^+ homeostasis. II. The effect of axonal stimulation, cholinergic agents and transport inhibition on the resistance in series with the axon membrane. *Neuroscience* **53**(3) 961-969 (1988)

5. Lieberman, E.M. and S. Hassan. Studies on axon-glia interactions and periaxonal K^+ homeostasis. III. The effect of anisosmotic media and potassium on the relationship between the resistance in series with the axon membrane and glial cell volume. *Neuroscience* 53:(3) 971-981 (1988)
6. Grossfeld, R., M.A. Klinge, E.M. Lieberman and L.C. Stewart. Axon-glia transfer of a protein and a carbohydrate. *Glia* 1:292-300 (1988)
7. Abbott, E.M., S. Hassan and E.M. Lieberman. Evidence for glutamate as the mediator of axon-Schwann cell interactions in the squid. *J. Physiol.(Lond.)* 398: 63P (1988)
8. Pichon, Y., N.J. Abbott, E.M. Lieberman and Y. Larmet. Potassium homeostasis in the nervous system of cephalopods and crustaceans. in: *J. Physiol (Paris)*. *J. Physiol. (Paris)* 82: 346-356 (1987)
9. Lieberman, E.M., Abbott, N.J. and Hassan, S. Evidence that glutamate mediates axon to Schwann signaling in the squid. *Glia* 2(3):94-102 (1989)
10. Lieberman, E.M. and Sanzenbacher, E. Possible mechanisms for glutamate mediation of axon-to-Schwann cell interaction in the isolated giant axon of the squid. *J. Physiol. (Lond.)* 423: 90P (1990)
11. Lieberman, E.M., Pascarella, J., Brunder, D. and Hargittai, P.T. The effect of extracellular potassium on ouabain-sensitive consumption of high energy phosphate by crayfish giant axons: A study of the energy requirement for transport in the steady-state. *J. Neurochem.* 55:155-164 (1990)
12. Hargittai, P.T., Butt, A.M. and Lieberman, E.M. High potassium selective permeability and extracellular ion regulation in the glial perineurium (Blood-Brain barrier) of the crayfish. *Neuroscience* 38: 163-173 (1990)
13. Butt, A.M., Hargittai, P.T. and Lieberman, E.M. Calcium-dependent potassium permeability in the glial perineurium (Blood-Brain barrier) of the crayfish. *Neuroscience* 38: 175-185 (1990)
14. Hargittai, P.T. and Lieberman E.M. Axon-glia interactions in the crayfish: Glial cell oxygen consumption is tightly coupled to axon metabolism. *Glia* 4:417-423 (1991)
15. Hargittai, P.T., Youmans, S. and Lieberman E.M. Determination of the membrane potential of cultured mammalian Schwann cells and its sensitivity to potassium using a thiocarbocyanine fluorescent dye. *Glia* 4:611-616 (1991)
16. Lieberman, E.M. Role of glutamate in axon-Schwann cell signaling in the squid. *Annals of the New York Academy of Science* 663:448-457 (1992)
17. Lieberman, E.M. and Sanzenbacher, E. Mechanisms of glutamate activation of axon-to-Schwann cell signaling in the squid. *Neuroscience* 47:931-939 (1992)
18. Stewart, W.C., Hargittai, P.T. and Lieberman, E.M. Cholera Toxin increases Na^+/K^+ -ATPase in the RN22 Schwann cell line. *Ann. N.Y. Acad. Sci.*

Abstracts: (work not yet published)

1. McKinnon, B., Hargittai, P.T. and Lieberman E.M. Immunocytochemical localization of glutamine synthetase and glutaminase in the giant axon of crayfish. Transactions of the American Society for Neurochemistry 24:203 (1993)

Manuscripts: (submitted or in press)

1. Lieberman, E.M., Hargittai, P.T. and Grossfeld, R.M. Electrophysiological and metabolic interactions between axons and glia in crayfish and squid. Progress in Neurobiology (Invited Review, 118 mnsct pages, submitted May 1993)
2. Nakhoul, N.L., Abdalnour-Nakhoul, S., Khuri, R.N., Lieberman, E.M. and Hargittai, P.T. Intracellular pH regulation in rat Schwann cells. Glia (1993) Submitted.

Inventions and Subcontracts:

No inventions resulted from the research supported under this grant nor subcontracts entered into.

Accession for	
NTIS	DTIC
DTIC	DTIC
DTIC	DTIC
DTIC	DTIC
By	
Dated	
Availability	
Dist	Av
A-1	

DTIC QUALITY INSPECTED 2